



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Bile acids in drug induced liver injury: Key players and surrogate markers

Schadt, Heiko S ; Wolf, Armin ; Pognan, Francois ; Chibout, Salah-Dine ; Merz, Michael ;
Kullak-Ublick, Gerd A

Abstract: Bile acid research has gained great momentum since the role of bile acids as key signaling molecules in the enterohepatic circulation was discovered. Their physiological function in regulating their own homeostasis, as well as energy and lipid metabolism make them interesting targets for the pharmaceutical industry in the context of diseases such as bile acid induced diarrhea, bile acid induced cholestasis or nonalcoholic steatohepatitis. Changes in bile acid homeostasis are also linked to various types of drug-induced liver injury (DILI). However, the key question whether bile acids are surrogate markers for monitoring DILI or key pathogenic players in the onset and progression of DILI is under intense investigation. The purpose of this review is to summarize the different facets of bile acids in the context of normal physiology, hereditary defects of bile acid transport and DILI.

DOI: <https://doi.org/10.1016/j.clinre.2015.12.017>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-128518>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Schadt, Heiko S; Wolf, Armin; Pognan, Francois; Chibout, Salah-Dine; Merz, Michael; Kullak-Ublick, Gerd A (2016). Bile acids in drug induced liver injury: Key players and surrogate markers. *Clinics and Research in Hepatology and Gastroenterology*, 40(3):257-266.

DOI: <https://doi.org/10.1016/j.clinre.2015.12.017>

Bile acids in drug induced liver injury: key players and surrogate markers

Heiko S. Schadt¹, Armin Wolf¹, Francois Pognan¹, Salah-Dine Chibout¹, Michael Merz¹, Gerd A. Kullak-Ublick^{2,3}

¹ Novartis Pharma AG, Preclinical Safety, Discovery and Investigative Safety, Basel, Switzerland

² Drug Safety and Epidemiology, Novartis Pharma AG, Basel, Switzerland

³ Department of Clinical Pharmacology and Toxicology, University Hospital Zurich, University of Zurich, Switzerland

Abstract

Bile acid research has gained great momentum since the role of bile acids as key signaling molecules in the enterohepatic circulation was discovered. Their physiological function in regulating their own homeostasis, as well as energy and lipid metabolism make them interesting targets for the pharmaceutical industry in the context of diseases such as bile acid induced diarrhea, bile acid induced cholestasis or nonalcoholic steatohepatitis. Changes in bile acid homeostasis are also linked to various types of drug-induced liver injury (DILI). However, the key question whether bile acids are surrogate markers for monitoring DILI or key pathogenic players in the onset and progression of DILI is under intense investigation. The purpose of this review is to summarize the different facets of bile acids in the context of normal physiology, hereditary defects of bile acid transport and DILI.

Introduction

The liver is highly exposed to drugs and xenobiotics, maintains the metabolic homeostasis of the body, and is responsible for bile formation and biliary secretion. Thus, it is not surprising that the liver is a major target organ of adverse drug reactions. Drug-induced liver injury can be classified as drug induced cholestasis, drug induced hepatitis and mixed type liver injury [1] and has been the most frequent single cause of safety-related drug marketing withdrawals for the past 50 years [2, 3]. DILI can resemble almost any form of acute or chronic liver disease (*e.g.* bland cholestasis, acute cholestatic hepatitis, acute hepatic

necrosis, acute viral hepatitis-like syndrome, immunoallergic hepatitis, cholestatic hepatitis due to medications, autoimmune hepatitis-like injury, sinusoidal obstruction syndrome, microvesicular steatosis with lactic acidosis and hepatic failure, fatty liver disease, chronic hepatitis, vanishing bile duct syndrome), thus necessitating standardized definitions and diagnostic criteria for DILI [4].

Bile acid biosynthesis and conjugation

Bile acids (BAs) are synthesized in the liver from cholesterol [5]. The classical (neutral) pathway is initiated by CYP7A1 catalyzed 7 α -hydroxylation, which represents the rate-limiting step in BA biosynthesis, leading to the formation of the two primary BAs cholic acid (CA) and chenodeoxycholic acid (CDCA). CA biosynthesis involves an additional hydroxylation at steroid ring position 12, catalyzed by CYP8B1, which controls the ratio of CA to CDCA formation. The side-chain is finally oxidized and cleaved by CYP27A1 [6]. This classical pathway produces at least 75 % of total BAs [7]. An alternate (acidic) pathway exists that is initiated by CYP27A1 and CYP7B1, leading to the formation of CDCA only [8, 9] [Figure 1]. In rodents, CDCA can be hydroxylated to α -muricholic acid (α MCA) and further epimerized to β MCA, giving rise to a pool of more polar BAs as compared to human [10]. Within the hepatocyte (and to a lesser extent in cholangiocytes), BAs are subjected to conjugation with glycine or taurine, which is catalyzed by the enzymes bile acid:CoA synthase (BACS) and bile acid:amino acid transferase (BAT) [11]. Dogs are predominant taurine-conjugators, rodents can form taurine and glycine conjugates, and non-human primates and humans show a preference for glycine conjugation. It has been shown that human BAT catalyzes BA conjugation with both glycine and taurine, and the extent to which these two conjugates are formed is determined by substrate availability [12]. This may also apply to other species with mixed taurine and glycine conjugation of BAs [13]. Other conjugation reactions such as sulfation and glucuronidation may occur, but are considered less relevant for the elimination of BAs under normal physiological conditions [14-20].

Transport of bile acids

After conjugation, BAs are excreted out of the hepatocyte across the canalicular membrane into the bile canaliculi, a process driven mainly by the bile salt export pump BSEP. This transport is ATP-dependent to facilitate BA excretion against a huge concentration gradient [21]. The relative rank order of intrinsic clearance for human BSEP is taurochenodeoxycholic acid (TCDCA) > taurocholic acid (TCA) > tauroursodeoxycholic acid (TUDCA) > glycocholic acid (GCA) [22-24]. In addition, multidrug resistance-associated protein 2 (MRP2) mediates the transport of divalent BA (*e.g.* sulfated and glucuronidated BA conjugated with taurine or glycine), conjugated bilirubin and glutathione-conjugates into bile [25]. P-glycoprotein (MDR1) appears to be capable to transport BAs across liver canalicular membranes to a certain degree [26], as shown in mice. Alternatively, basolateral export of BAs from hepatocytes and cholangiocytes back into the systemic circulation is mediated by multidrug resistance proteins MRP3/MRP4 and organic solute transporters OST- α /OST- β [21, 25, 27-29]. Cholangiocytes, similar to gallbladder epithelial cells, proximal renal convoluted tubule cells and ileal enterocytes, express the sodium-dependent BA transporter ASBT at their apical membrane domain, thereby allowing for cholehepatic shunting of conjugated BAs back to hepatocytes [30].

Cholangiocytes

The amphipathic nature of bile salts drives the solubilisation of phosphatidylcholine and cholesterol in bile canaliculi by forming mixed micelles, thus protecting cholangiocytes from the detergent properties of the bile salts. During its passage through the biliary tree, the primary hepatic bile is further modified by organic anion and electrolyte transport proteins expressed at the surface of biliary epithelial cells [22]. Cholangiocytes secrete significant amounts of bicarbonate, which raises the local pH at the apical membrane. Beuers *et al.* proposed a concept termed the “biliary bicarbonate umbrella” [31] and hypothesized that biliary bicarbonate secretion near the apical surface of hepatocytes and cholangiocytes maintains an alkaline milieu that keeps the predominantly glycine-conjugated BAs in man in their salt form and therefore in a more polar state. Polarity prevents molecules from passive diffusion through

biological membranes, which could therefore protect the integrity of cholangiocytes and periportal hepatocytes from the toxicity of highly concentrated bile salts. Dysregulation or functional impairment of this biliary bicarbonate umbrella may consequently increase the vulnerability of biliary epithelial cells towards biliary bile salts, and therefore contribute to various forms of chronic cholangiopathies/cholestatic liver diseases [31]. Supporting evidence comes from the anion exchanger AE2, which is expressed on the apical membrane of cholangiocytes and secretes bicarbonate into the biliary lumen in exchange for chloride. AE2 deficient knockout mice develop features resembling primary biliary cirrhosis [32]. Human genetic variants in this gene have been shown to influence susceptibility and severity of primary biliary cirrhosis [33]. In vitro experiments with human cholangiocytes in the presence of millimolar concentrations of CDCA and its glycine- and taurine conjugates confirmed the pH- and AE2-dependent uptake and cytotoxicity of BAs. In addition, these experiments suggested a potential protective role of the intact biliary glycocalyx against pH-dependent bile salt toxicity [34, 35].

Enterohepatic cycling of bile acids

In humans, dog and mice, bile is stored in the gallbladder (which is absent in rats) before it is released upon the action of the hormone cholecystokinin (CCK) into the small intestine to facilitate lipid digestion. BA can be deconjugated and further modified (e.g. dehydroxylation) by the intestinal microflora, giving rise to secondary, more hydrophobic BAs. Primary CA derived BAs (CA, GCA, TCA) are converted to the secondary BA deoxycholic acid (DCA), whereas primary CDCA derived BAs (CDCA, TCDCA, glycochenodeoxycholic acid (GCDCA)) form the secondary BA lithocholic acid (LCA). In rats, the primary BA β MCA is transformed to the secondary BA ω MCA by various anaerobic bacteria in a cooperative way [36]. BAs in the intestine are reabsorbed either by passive diffusion (e.g. DCA) or by active sodium-dependent transport *via* the ASBT transporter into enterocytes [37]. BAs escaping intestinal absorption are excreted *via* feces accounting for about 10 to 20 % loss per day [7], fecal BAs being mainly secondary BAs. From enterocytes, BAs enter the portal circulation via the OST- α /OST- β transporter heterodimer and are taken up into the liver across the hepatocyte sinusoidal membrane by the sodium-

(Na⁺)-taurocholate cotransporting polypeptide (NTCP) or by organic anion transporting polypeptides (OATPs). The reabsorbed BAs mix with *de novo* synthesized BA inside hepatocytes and can be re-conjugated before entering the next enterohepatic cycle [38].

Regulation of bile acid homeostasis

The enterohepatic FXR-FGF15/FGF19-FGFR4/βKlotho pathway assigned a new endocrine role to BAs [39]. Within enterocytes BAs bind to the farnesoid X nuclear receptor FXR and activate the transcription of fibroblast growth factor FGF15/19. FGF19 in humans, and FGF15 in rodents, is released into the portal blood from where it reaches the liver and binds to and activates the membrane-bound FGFR4/βKlotho receptor complex [40]. Activation of this receptor tyrosine kinase triggers a complex intracellular signaling cascade that finally results in the suppression of CYP7A1 - the key regulatory enzyme of *de novo* BA synthesis in hepatocytes. By this enterohepatic feedback circuit BAs regulate their own biosynthesis in an endocrine manner [41]. Additional feedback-mechanisms exist in the liver for controlling BA homeostasis, both at the transcriptional and posttranscriptional levels. FXR is also expressed in hepatocytes and regulates hepatic BA biosynthesis and transport proteins that are involved in hepatocellular uptake (NTCP, OATP) and excretion (BSEP, MRP2, OST-α/OST-β, MDR1). Activation of FXR by BAs leads to suppression of both CYP7A1, CYP8B1 and the hepatic influx transporters NTCP and OATP (human only), and to an upregulation of the hepatic efflux transporters BSEP, MRP2, OST-α/OST-β, thus preventing accumulation of cytotoxic BAs within the hepatocytes under normal physiologic conditions [38, 42-45]. The relative potency of individual BAs to activate FXR was determined as CDCA>DCA>LCA. The corresponding glycine- and taurine conjugates were also shown to activate FXR when BA transporters were co-expressed in cellular *in vitro* test systems. Free and conjugated CA, ursodeoxycholic acid (UDCA) and α/βMCA had only minimal or no effect on FXR [46, 47].

BA in health and disease

The concept that BAs could be key players in DILI is derived from the fact that BAs are amphiphilic molecules with strong emulsifying detergent properties that may impair cell membrane integrity and thus exert cytotoxic effects. The liver is exposed to extremely high concentrations of BAs (canaliculus and biliary ductules: 20 to 50 mM, gallbladder: ≤ 300 mM), which makes this organ particularly vulnerable to BA-induced toxicity [48].

The lipophilicity of individual BAs is believed to be a determinant of cytotoxicity, since it is the main driver for passive diffusion of BAs through biological membranes. In addition, the pH of the surrounding environment together with the pKs of the individual BA determines whether the molecule exists in its uncharged, protonated or salt form, which makes a significant difference for its polarity and capability to penetrate membranes [49, 50]. Unconjugated DCA, LCA, CDCA, but also conjugated GCDCA and TCDCA were reported to impair the functionality of mitochondria isolated from rat liver, which may in part mediate cytotoxicity [51]. Structure-activity analysis of BAs showed that the cytotoxicity of naturally occurring BAs can be ranked according to $DCA \geq LCA \geq CDCA \gg CA$ based on assessment in an oesophageal (HET-1A) and hepatic carcinoma (Huh7) cell line, and overall this rank order correlates with lipophilicity. However, the respective taurine and glycine conjugates had no cytotoxic effect on HET-1A cells, which lack BA transporters [52]. Conjugated BAs are fully ionized at physiological pH and are therefore considered membrane impermeable without active transport. Thus, intracellular toxicity mediated by conjugated BAs only occurs when transporters facilitate their uptake into the cell [48] or if their efflux is inhibited leading to intracellular accumulation. It consequently appears appropriate to consider transporter-mediated flux of conjugated BAs into different compartments along with potential interconversion reactions that may affect their cytotoxicity profile when discussing the role of BAs in DILI. Figure 2A illustrates the differences in the BA profiles in plasma, bile, feces and urine of dogs as determined by LC-MS/MS in our laboratory.

It is well known that different species show different BA pool compositions (Figure 2B), which adds additional complexity to the interpretation of preclinical data in the context of human safety assessment during drug development. Rodents, for example, have a higher portion of polar BAs due to the synthesis of a higher proportion of hydroxylated MCAs, which makes their BA pool generally less cytotoxic. Interestingly, genetically modified mouse models susceptible towards developing progressive familial intrahepatic cholestasis (PFIC) display only a mild phenotype compared to human patients. However, when the BA pool of these PFIC susceptible mice was “humanized” by liver-specific disruption of the cytochrome P450 reductase gene resulting in reduced CYP450 activity and cholic acid supplementation, a more hydrophobic BA pool was established and mice rapidly developed pronounced cholestasis with histologic liver injury [53]. These data might therefore indicate that BA pool composition and changes thereof could be a critical factor for the susceptibility to liver injury.

The mutual effects of BAs on the intestinal microflora and *vice versa* have repeatedly been reported [54]. The composition of the gut microbiota and their metabolic capacity strongly affects the composition of the BA pool, notably its hydrophobicity, cytotoxicity and size [55]. Treatment of mice with probiotics resulted in enhanced deconjugation and fecal excretion of BAs, affecting ileal BA absorption and subsequent stimulation of hepatic *de novo* BA synthesis via repression of the enterohepatic FXR-FGF15/FGF19-FGFR4 pathway [56]. Hara and coworkers reported that obesity induced changes of the gut microbiota, leading to increased levels of the secondary BA DCA in the enterohepatic circulation. DCA, being a relatively cytotoxic BA, provoked DNA damage and cellular senescence in hepatic stellate cells, which consequently released inflammatory and tumor-promoting factors in the liver, facilitating hepatocellular carcinoma development in mice previously exposed to a chemical carcinogen [57]. Drugs that impair the intestinal microflora can therefore indirectly affect the liver by altering the BA profile and BA induced cellular effects [58, 59].

Various liver diseases are associated with impaired BA homeostasis. Genetic variations in the genes for FXR, BSEP and MDR3 confer susceptibility to intrahepatic cholestasis of pregnancy (ICP) - a transient

cholestasis in pregnant women – in association with elevated circulating levels of estrogens and progesterone [22]. Estrogen and progesterone metabolites, *e.g.* estradiol-17 β -glucuronide, were shown to trans-inhibit BSEP after its secretion into bile canaliculi by MRP2 [60, 61]. Increased serum levels of sulfated progesterone metabolites were reported to affect FXR functionality in addition to inhibiting the hepatocellular BA uptake and efflux transporters NTCP and BSEP [62-64], thus mechanistically linking the cholestatic syndrome in susceptible women to the hormonal changes arising during pregnancy. Consistently, drug therapy with oral contraceptives and anabolic steroids such as 17 α -alkylated androgens were also associated with cholestasis, depending on the individual susceptibility of the recipient [65].

Progressive familial intrahepatic cholestasis (PFIC) is an inherited early-onset cholestatic syndrome with progression to end-stage liver disease caused by mutations in canalicular transporter genes. Less severe forms of cholestasis such as benign recurrent intrahepatic cholestasis (BRIC) are linked to mutations in BSEP or MDR3, likely impairing expression or function of the respective transport protein [22]. Treatment of ICP patients with the secondary BA UDCA was reported to reduce plasma concentrations of sulfated steroid metabolites [66] and exert other beneficial effects [67]. These and various other examples illustrate the importance of intact BA homeostasis for the integrity of the liver.

Bile acids as key players in DILI:

Consistent with the role of BA transporter impairment in various liver diseases, inhibition of BSEP by drugs or their metabolites is considered an important mechanism of drug induced cholestasis and has been reported for cyclosporine A, rifampicin, bosentan, troglitazone and various other compounds [68].

While drug induced cholestasis without hepatitis can be transient and is commonly less severe for the patient, severe irreversible drug-induced liver failure can be fatal or require organ transplantation. Troglitazone (TGZ) is an anti-diabetic drug that was withdrawn from the market due to acute severe idiosyncratic drug-induced liver injury in some patients [69-71]. Clinical observations and *in vivo* rat studies suggested, at least in part, a cholestatic mechanism for TGZ-induced hepatotoxicity, in addition to

a role of a reactive quinone metabolite [72, 73]. Both in human and in rat, TGZ is efficiently metabolized to the major metabolite TGZ-sulfate, which has a high potential to competitively inhibit BSEP and accumulate in the hepatocytes (apparent K_i for rat BSEP inhibition were determined as 1.3 μM for TGZ and 0.23 μM for TGZ-sulfate) [74]. Rat studies with TGZ demonstrated significantly reduced bile flow and a rapid and dose-dependent increase in plasma BA concentrations [74, 75]. A mechanism leading to cytotoxic intracellular bile salt accumulation due to BSEP inhibition was therefore also discussed for humans. Despite the disturbances in BA homeostasis, rat models failed to predict TGZ-induced hepatotoxicity, potentially due to interspecies differences in BA composition and the less cytotoxic BA pool in rodents [76].

Jemnitz *et al.* investigated the effect of cholestatic drugs on TCA transport in human and rat sandwich-cultured hepatocytes and found significant differences in TCA elimination: in human hepatocytes, equal fractions of TCA were eliminated via basolateral (likely by MRP3/4 efflux pumps) and canalicular (mainly BSEP) efflux (34.8% and 34.4 %, respectively) with a similar portion remaining within the cell (30.5%). In contrast, basolateral transport (71.7%) was dominant in rat hepatocytes and intracellular TCA accumulation was low (6.9%). Concordantly, inhibition of BSEP by TGZ treatment resulted in a significantly higher relative increase in intracellular TCA concentrations in human *versus* rat hepatocytes [77]. The higher rate of basolateral BA efflux in the rat in combination with the lower intracellular concentrations and the lower toxicity of the rodent BA pool could have a protective role in the rat and explain species differences in response to cholestatic or hepatotoxic drugs.

Recently, a system pharmacology model was described that could predict species differences in TGZ-induced hepatotoxicity due to altered BA homeostasis [76]. The model simulated a human population, taking into account drug/metabolite disposition, BA physiology/pathophysiology (transporter inhibition), hepatocyte life cycle and liver injury biomarkers and predicted TGZ-induced delayed increases in serum alanine transaminase and bilirubin levels. Such physiologically based pharmacokinetic models could be of great value for assessing adverse outcome effects due to impairment of BA homeostasis. Implementation

of additional compensatory regulatory networks, *e.g.* FXR-mediated transcriptional up- or downregulation of hepatic BA transporters and biosynthesis enzymes, or the endocrine FXR-FGF19/FGFR4 pathway, could add another valuable dimension.

The drug bosentan, an endothelin receptor antagonist approved for pulmonary arterial hypertension, contains a black-box warning for hepatotoxicity. Similarly to TGZ, a mechanism involving inhibition of BSEP with subsequent intracellular accumulation of cytotoxic BAs and induction of liver cell damage was discussed [78]. Despite the inhibitory potential on rat BSEP leading to elevated serum BAs, bosentan did not induce hepatotoxicity in rats. Lesli *et al.* showed that bosentan differentially inhibits both rat and human NTCP in addition to BSEP. The authors concluded that the rat might be less sensitive to hepatotoxicity since bosentan showed stronger inhibitory potential towards rat NTCP compared to human, resulting in less intrahepatocyte accumulation of BAs in rats [79]. By applying a mechanistic model of DILI Woodhead *et al.* simulated whether - in response to bosentan - BA accumulation can reach toxic levels, both in a human and in a rat population, or not. Although the incidence of bosentan toxicity was underestimated, the system predicted a mild hepatocellular ATP decline and serum ALT elevation in human but not in rats, which was accounted for by the less toxic BA pool in rats [80].

A recent case report described two Japanese patients who had initially received bosentan monotherapy and experienced liver toxicity. After dose reduction or discontinuation, liver function was normalized and patients received retreatment with a combination of bosentan and UDCA. Subsequent liver function tests did not show any abnormalities after the combined therapy [81], indicating compensatory effects by the treatment with UDCA.

Impairment of BA homeostasis by a single insult, *e.g.* inhibition of a single hepatic BA transporter, might be compensated by activation of regulatory feedback pathways involving nuclear receptors (*e.g.* FXR), leading to up- or down-regulation of hepatic BA transport systems. Additionally, alternative BA efflux systems or detoxifying BA conjugation reactions could prevent intracellular accumulation of cytotoxic BAs. However, additional insults by a drug impairing the integrity of multiple compensatory pathways

could exceed the protective capacity of the liver and become critical for liver function. Glucuronidation of BAs, for example, appears to be of little relevance under normal physiologic conditions [82]. However, it was reported that glucuronidation may become more important under severe cholestatic conditions, since it allows the conversion of cytotoxic BAs into non-toxic glucuronides, which also facilitates MRP3-mediated alternative efflux from hepatocytes into the blood for urinary elimination [16, 83-85]. Sulfation of BAs by sulfotransferase (*e.g.* SULT2A1) is another important mechanism to detoxify BAs. Sulfated BAs exhibit decreased toxicity and enhanced fecal and urinary excretion since they cannot be reabsorbed by the intestine, as exemplified by LCA sulfates. Sulfation of BAs is increased during cholestatic disease and might therefore play an important role under pathologic conditions [86].

Although there are various indications that BAs might be key players in the onset of DILI, the initiation and progression of liver injury is usually multifactorial. Aleo *et al.*, for example, assessed various compounds with Most-DILI-, Less-DILI- and No-DILI-concern and found that drugs with dual potency as mitochondrial and BSEP inhibitors were highly associated with more severe human DILI and more restrictive product safety labeling related to liver injury [87]. In the case of TGZ, reactive metabolite formation and mitochondrial dysfunction are considered to be critical steps in DILI pathogenesis in addition to inhibition of BSEP [73, 74, 88]. Inflammatory processes significantly contribute to DILI, which may often be triggered by reactive metabolite mediated hapten formation. However, BAs were also shown to induce inflammatory genes in hepatocytes and in mouse liver after bile duct ligation [89], which may promote hepatic inflammation during cholestasis and contribute to DILI. Besides the potential contribution of impaired BA homeostasis, mechanisms such as mitochondrial dysfunction, cytotoxicity, reactive metabolite formation, immune-mediated hypersensitivity reactions, dose and lipophilicity of a drug itself and combinations thereof [90, 91] seem to be important factors in the onset of acute liver failure in combination with individual risk factors.

Bile acids as surrogate markers

Circulating total BAs are well-established clinical markers to detect liver injury and reflect a disturbance of normal hepatic excretory function. Changes in total BAs are usually indicative for hepatocellular and/or hepatobiliary injury. With the availability of new analytical multiplexing methods based on LC-MS/MS and GC/MS, circulating BA profiles are currently being evaluated as biomarkers for hepatotoxicity [92]. In an attempt to identify potential biomarkers for improved DILI detection by untargeted metabolomics, significant elevations of a panel of BAs in plasma and urine of rats treated with different types of hepatotoxins were found [93]. Other groups reported changes in metabolites derived from energy [94, 95], urea and BA metabolism in urine and blood of mice or rats treated with a single dose of acetaminophen (APAP), which strongly correlated with the development of hepatic necrosis [96]. Luo *et al.* evaluated by a targeted LC-MS/MS approach CA, GCA and TCA as potential biomarkers of liver injury in rodent models [97]. The authors reported that animals with histopathological signs of hepatocellular necrosis showed increases in all three BAs in serum, whereas animals with signs of bile duct hyperplasia displayed increases in only the conjugated forms GCA and TCA. They suggested that individual serum BAs might be a potential diagnostic marker to differentiate between different forms of liver injury [97] and characterized individual BA profiles in patients with liver injury and healthy volunteers [98]. Their data showed that total and individual BAs were strongly increased in samples from hepatic injury subjects with the exception of the secondary BA DCA. In addition, conjugated BAs were higher than unconjugated BAs in the patient group. The portion of free and conjugated CA in serum was also higher in subjects with liver injury compared to healthy controls. The changes induced by APAP with respect to individual BAs in urine and blood based on clinical and non-clinical data and their potential use as biomarkers of liver injury were recently reported [99].

While alterations in BA profiles can be the consequence of hepatotoxicity, functional inhibition of BA transporters *in vivo* may also lead to shifts in BA composition in the circulation without being the consequence of cellular damage of the liver. In an exploratory study with male Sprague-Dawley rats

treated with 20 mg/kg/day cyclosporine A (CsA) by oral gavage for 7 days, we found significant changes in individual plasma BA components. The unconjugated primary BAs CA, CDCA, and α/β MCA, which accounted for approximately 60% of the total BAs in control plasma, were significantly increased with CsA. While DCA plasma levels were not affected by the treatment, the unconjugated secondary BAs ω MCA, γ MCA and UDCA tended to be increased in absolute concentrations. LCA, though present only in traces in plasma, was decreased. Both primary and secondary taurine-conjugated BAs were generally less abundant in treated animals when compared to controls. The sum of all individual BAs measured increased approximately 3-fold with CsA, which was mainly driven by the increase of the unconjugated primary BAs CA, CDCA and α MCA (Figure 2C). This complex pattern of changes in circulating BAs observed with CsA in rats is likely the consequence of inhibition of multiple hepatic transporters. CsA has the potential to inhibit both rat influx and efflux transporters in the lower micromolar range as determined in primary cultured rat hepatocytes and membrane vesicles, respectively (NTCP IC_{50} = 8.63 μ M, OATPs IC_{50} = 4.55 μ M, BSEP IC_{50} = 1.75 μ M, MRP2 IC_{50} = 9.2 μ M) [100]. Although the individual contribution of these transporters to CsA-induced changes in BA profiles is difficult to assign, it seems that not all BAs were affected in the same way. To further shed light into the complex system of BA homeostasis, it would be advantageous to systematically generate structure-activity relations of BAs for their various BA transporters. Such data could be valuable to better simulate potential BA-mediated adverse effects in humans, taking the different species' BA profiles into account.

The IMI SAFE-T consortium, a public-private partnership between industry, health authorities and academia, is currently evaluating BA profiles as potential markers for drug-induced BA transporter inhibition and liver injury [101]. The complexity of this endeavor lies in the fact that different mechanisms can lead to changes in BA profiles with overlapping effects. Many drugs inhibit more than one BA transporter and/or have effects on BA biosynthesis. In addition, the transition from BAs as early functional markers to assess BA transporter inhibition to surrogate biomarkers for liver injury may be difficult to differentiate based on thresholds. Preclinical studies can be very helpful since they allow an optimal study design with controlled conditions and appropriate sampling. However, the differences between species in

BA composition and physiology need to be considered for proper data interpretation and subsequent translation or extrapolation to humans. Human biosamples with a clearly defined pathology are mandatory for validating circulating individual BAs as biomarkers for monitoring liver integrity.

Summary and conclusion

Accumulation of intrahepatic cytotoxic BAs due to drug-induced disturbances of BA homeostasis appears to be an important contributing factor to the onset of DILI in susceptible individuals in combination with other key adverse events impacting liver physiology, *e.g.* mitochondrial dysfunction, reactive metabolite formation and immune-mediated responses. Circulating individual BAs' concentrations might become promising sensitive and selective early markers to monitor functional impairment of the liver, allowing a prognostic assessment before irreversible damage occurs.

Acknowledgements

We thank Marianne Schwald and Audrey Fischer for excellent technical support with LC-MS/MS based multiplexing bile acid profile analysis.

Figure legends

Fig. 1: Schematic overview of hepatic BA biosynthesis, conjugation, transport and enterohepatic cycling. Regulatory pathways of BA homeostasis, *e.g.* the endocrine FXR-FGF15/19-FGFR4 pathway and intrahepatic regulation of BA transporters by FXR, are shown. The rodent specific MCAs are depicted in *italic*

Fig. 2: A: BA profiles in dog plasma, bile, feces and urine. B: Differences in BA composition in plasma of mouse, rat, monkey and human. C: Changes in individual plasma BA concentrations in rats treated with 20 mg/kg/day Cyclosporine A for 7 days. Levels of statistical significance *versus* controls were calculated using a student's *t* test: * $p > 0.05$; ** $p > 0.01$; *** $p > 0.001$. Unconj.1stBA: sum of unconjugated primary BAs. T-1stBA: sum of taurine conjugated primary BAs. T-2ndBA: sum of taurine conjugated secondary BAs. All data were obtained in-house by a multiplexing LC-MS/MS method enabling the analysis of 17 BAs.

References

1. Standardization of definitions and criteria of causality assessment of adverse drug reactions. Drug-induced liver disorders: report of an international consensus meeting. *Int J Clin Pharmacol Ther Toxicol.* 1990;28(8):317-22.
2. Lasser KE, Allen PD, Woolhandler SJ, Himmelstein DU, Wolfe SM, Bor DH. Timing of new black box warnings and withdrawals for prescription medications. *JAMA.* 2002;287(17):2215-20.
3. Temple RJ, Himmel MH. Safety of newly approved drugs: implications for prescribing. *JAMA.* 2002;287(17):2273-5.
4. Fontana RJ, Seeff LB, Andrade RJ, Bjornsson E, Day CP, Serrano J, et al. Standardization of nomenclature and causality assessment in drug-induced liver injury: summary of a clinical research workshop. *Hepatology.* 2010;52(2):730-42.
5. Myant NB, Mitropoulos KA. Cholesterol 7 α -hydroxylase. *J Lipid Res.* 1977;18(2):135-53.
6. Chiang JY. Bile acid regulation of hepatic physiology: III. Bile acids and nuclear receptors. *Am J Physiol Gastrointest Liver Physiol.* 2003;284(3):G349-56.
7. Thomas C, Pellicciari R, Pruzanski M, Auwerx J, Schoonjans K. Targeting bile-acid signalling for metabolic diseases. *Nat Rev Drug Discov.* 2008;7(8):678-93.

8. Russell DW. The enzymes, regulation, and genetics of bile acid synthesis. *Annu Rev Biochem.* 2003;72:137-74.
9. Russell DW. Fifty years of advances in bile acid synthesis and metabolism. *J Lipid Res.* 2009;50 Suppl:S120-5.
10. Zhang Y, Klaassen CD. Effects of feeding bile acids and a bile acid sequestrant on hepatic bile acid composition in mice. *J Lipid Res.* 2010;51(11):3230-42.
11. Chiang JY. Bile acids: regulation of synthesis. *J Lipid Res.* 2009;50(10):1955-66.
12. Falany CN, Johnson MR, Barnes S, Diasio RB. Glycine and taurine conjugation of bile acids by a single enzyme. Molecular cloning and expression of human liver bile acid CoA:amino acid N-acyltransferase. *J Biol Chem.* 1994;269(30):19375-9.
13. Sturman JA, Messing JM, Rossi SS, Hofmann AF, Neuringer M. Tissue taurine content, activity of taurine synthesis enzymes and conjugated bile acid composition of taurine-deprived and taurine-supplemented rhesus monkey infants at 6 and 12 mo of age. *J Nutr.* 1991;121(6):854-62.
14. Hofmann AF, Hagey LR. Key discoveries in bile acid chemistry and biology and their clinical applications: history of the last eight decades. *J Lipid Res.* 2014;55(8):1553-95.
15. Matern S, Matern H, Farthmann EH, Gerok W. Hepatic and extrahepatic glucuronidation of bile acids in man. Characterization of bile acid uridine 5'-diphosphate-glucuronosyltransferase in hepatic, renal, and intestinal microsomes. *J Clin Invest.* 1984;74(2):402-10.
16. Trottier J, Verreault M, Grepper S, Monte D, Belanger J, Kaeding J, et al. Human UDP-glucuronosyltransferase (UGT)1A3 enzyme conjugates chenodeoxycholic acid in the liver. *Hepatology.* 2006;44(5):1158-70.
17. Takikawa H, Beppu T, Seyama Y. Urinary concentrations of bile acid glucuronides and sulfates in hepatobiliary diseases. *Gastroenterol Jpn.* 1984;19(2):104-9.
18. Radomska-Pyrek A, Zimniak P, Chari M, Golunski E, Lester R, St Pyrek J. Glucuronides of monohydroxylated bile acids: specificity of microsomal glucuronyltransferase for the glucuronidation site, C-3 configuration, and side chain length. *J Lipid Res.* 1986;27(1):89-101.
19. Shattuck KE, Radomska-Pyrek A, Zimniak P, Adcock EW, Lester R, St Pyrek J. Metabolism of 24-norlithocholic acid in the rat: formation of hydroxyl- and carboxyl-linked glucuronides and effect on bile flow. *Hepatology.* 1986;6(5):869-73.
20. Bock KW. Roles of human UDP-glucuronosyltransferases in clearance and homeostasis of endogenous substrates, and functional implications. *Biochem Pharmacol.* 2015;96(2):77-82.
21. Dawson PA, Lan T, Rao A. Bile acid transporters. *J Lipid Res.* 2009;50(12):2340-57.
22. Kullak-Ublick GA, Stieger B, Meier PJ. Enterohepatic bile salt transporters in normal physiology and liver disease. *Gastroenterology.* 2004;126(1):322-42.
23. Byrne JA, Strautnieks SS, Mieli-Vergani G, Higgins CF, Linton KJ, Thompson RJ. The human bile salt export pump: characterization of substrate specificity and identification of inhibitors. *Gastroenterology.* 2002;123(5):1649-58.
24. Noe J, Stieger B, Meier PJ. Functional expression of the canalicular bile salt export pump of human liver. *Gastroenterology.* 2002;123(5):1659-66.
25. Meier PJ, Stieger B. Bile salt transporters. *Annu Rev Physiol.* 2002;64:635-61.
26. Lam P, Wang R, Ling V. Bile acid transport in sister of P-glycoprotein (ABCB11) knockout mice. *Biochemistry.* 2005;44(37):12598-605.
27. Li N, Cui Z, Fang F, Lee JY, Ballatori N. Heterodimerization, trafficking and membrane topology of the two proteins, Ost alpha and Ost beta, that constitute the organic solute and steroid transporter. *Biochem J.* 2007;407(3):363-72.
28. Seward DJ, Koh AS, Boyer JL, Ballatori N. Functional complementation between a novel mammalian polygenic transport complex and an evolutionarily ancient organic solute transporter, OSTalpha-OSTbeta. *J Biol Chem.* 2003;278(30):27473-82.

29. Wang W, Seward DJ, Li L, Boyer JL, Ballatori N. Expression cloning of two genes that together mediate organic solute and steroid transport in the liver of a marine vertebrate. *Proc Natl Acad Sci U S A*. 2001;98(16):9431-6.
30. Lazaridis KN, Pham L, Tietz P, Marinelli RA, deGroen PC, Levine S, et al. Rat cholangiocytes absorb bile acids at their apical domain via the ileal sodium-dependent bile acid transporter. *J Clin Invest*. 1997;100(11):2714-21.
31. Beuers U, Hohenester S, de Buy Wenniger LJ, Kremer AE, Jansen PL, Elferink RP. The biliary HCO₃(-) umbrella: a unifying hypothesis on pathogenetic and therapeutic aspects of fibrosing cholangiopathies. *Hepatology*. 2010;52(4):1489-96.
32. Salas JT, Banales JM, Sarvide S, Recalde S, Ferrer A, Uriarte I, et al. Ae2a,b-deficient mice develop antimitochondrial antibodies and other features resembling primary biliary cirrhosis. *Gastroenterology*. 2008;134(5):1482-93.
33. Poupon R, Ping C, Chretien Y, Corpechot C, Chazouilleres O, Simon T, et al. Genetic factors of susceptibility and of severity in primary biliary cirrhosis. *J Hepatol*. 2008;49(6):1038-45.
34. Hohenester S, Wenniger LMdB, Paulusma CC, van Vliet SJ, Jefferson DM, Elferink RPO, et al. A biliary HCO₃- umbrella constitutes a protective mechanism against bile acid-induced injury in human cholangiocytes. *Hepatology*. 2012;55(1):173-83.
35. Maillette de Buy Wenniger LJ, Hohenester S, Maroni L, van Vliet SJ, Oude Elferink RP, Beuers U. The Cholangiocyte Glycocalyx Stabilizes the 'Biliary HCO₃ Umbrella': An Integrated Line of Defense against Toxic Bile Acids. *Dig Dis*. 2015;33(3):397-407.
36. Eysen H, De Pauw G, Stragier J, Verhulst A. Cooperative formation of omega-muricholic acid by intestinal microorganisms. *Appl Environ Microbiol*. 1983;45(1):141-7.
37. Dawson PA. Role of the intestinal bile acid transporters in bile acid and drug disposition. *Handb Exp Pharmacol*. 2011(201):169-203.
38. Kuipers F, Bloks VW, Groen AK. Beyond intestinal soap--bile acids in metabolic control. *Nat Rev Endocrinol*. 2014;10(8):488-98.
39. Inagaki T, Choi M, Moschetta A, Peng L, Cummins CL, McDonald JG, et al. Fibroblast growth factor 15 functions as an enterohepatic signal to regulate bile acid homeostasis. *Cell Metab*. 2005;2(4):217-25.
40. Lin BC, Wang M, Blackmore C, Desnoyers LR. Liver-specific activities of FGF19 require Klotho beta. *J Biol Chem*. 2007;282(37):27277-84.
41. Houten SM, Watanabe M, Auwerx J. Endocrine functions of bile acids. *EMBO J*. 2006;25(7):1419-25.
42. Kullak-Ublick GA. ABC transporter regulation by bile acids: where PXR meets FXR. *J Hepatol*. 2003;39(4):628-30.
43. Landrier JF, Eloranta JJ, Vavricka SR, Kullak-Ublick GA. The nuclear receptor for bile acids, FXR, transactivates human organic solute transporter-alpha and -beta genes. *Am J Physiol Gastrointest Liver Physiol*. 2006;290(3):G476-85.
44. Goodwin B, Jones SA, Price RR, Watson MA, McKee DD, Moore LB, et al. A regulatory cascade of the nuclear receptors FXR, SHP-1, and LXR-1 represses bile acid biosynthesis. *Mol Cell*. 2000;6(3):517-26.
45. Geier A, Wagner M, Dietrich CG, Trauner M. Principles of hepatic organic anion transporter regulation during cholestasis, inflammation and liver regeneration. *Biochim Biophys Acta*. 2007;1773(3):283-308.
46. Parks DJ, Blanchard SG, Bledsoe RK, Chandra G, Consler TG, Kliewer SA, et al. Bile acids: natural ligands for an orphan nuclear receptor. *Science*. 1999;284(5418):1365-8.
47. Makishima M, Okamoto AY, Repa JJ, Tu H, Learned RM, Luk A, et al. Identification of a nuclear receptor for bile acids. *Science*. 1999;284(5418):1362-5.
48. Hofmann AF. The continuing importance of bile acids in liver and intestinal disease. *Arch Intern Med*. 1999;159(22):2647-58.

49. Roda A, Grigolo B, Pellicciari R, Natalini B. Structure-activity relationship studies on natural and synthetic bile acid analogs. *Dig Dis Sci.* 1989;34(12 Suppl):24S-35S.
50. Roda A, Minutello A, Angellotti MA, Fini A. Bile acid structure-activity relationship: evaluation of bile acid lipophilicity using 1-octanol/water partition coefficient and reverse phase HPLC. *J Lipid Res.* 1990;31(8):1433-43.
51. Rolo AP, Oliveira PJ, Moreno AJ, Palmeira CM. Bile acids affect liver mitochondrial bioenergetics: possible relevance for cholestasis therapy. *Toxicol Sci.* 2000;57(1):177-85.
52. Sharma R, Majer F, Peta VK, Wang J, Keaveney R, Kelleher D, et al. Bile acid toxicity structure-activity relationships: correlations between cell viability and lipophilicity in a panel of new and known bile acids using an oesophageal cell line (HET-1A). *Bioorg Med Chem.* 2010;18(18):6886-95.
53. Kunne C, de Graaff M, Duijst S, de Waart DR, Oude Elferink RP, Paulusma CC. Hepatic cytochrome P450 deficiency in mouse models for intrahepatic cholestasis predispose to bile salt-induced cholestasis. *Lab Invest.* 2014;94(10):1103-13.
54. Devkota S, Chang EB. Interactions between Diet, Bile Acid Metabolism, Gut Microbiota, and Inflammatory Bowel Diseases. *Dig Dis.* 2015;33(3):351-6.
55. Ridlon JM, Kang DJ, Hylemon PB, Bajaj JS. Bile acids and the gut microbiome. *Curr Opin Gastroenterol.* 2014;30(3):332-8.
56. Degirolamo C, Rainaldi S, Bovenga F, Murzilli S, Moschetta A. Microbiota modification with probiotics induces hepatic bile acid synthesis via downregulation of the Fxr-Fgf15 axis in mice. *Cell Rep.* 2014;7(1):12-8.
57. Yoshimoto S, Loo TM, Atarashi K, Kanda H, Sato S, Oyadomari S, et al. Obesity-induced gut microbial metabolite promotes liver cancer through senescence secretome. *Nature.* 2013;499(7456):97-101.
58. Ridlon JM, Kang DJ, Hylemon PB, Bajaj JS. Gut microbiota, cirrhosis, and alcohol regulate bile acid metabolism in the gut. *Dig Dis.* 2015;33(3):338-45.
59. Bajaj JS, Cox LJ, Betrapally NS, Heuman DM, Schubert ML, Ratneswaran M, et al. Systems biology analysis of omeprazole therapy in cirrhosis demonstrates significant shifts in gut microbiota composition and function. *Am J Physiol Gastrointest Liver Physiol.* 2014;307(10):G951-7.
60. Stieger B, Fattinger K, Madon J, Kullak-Ublick GA, Meier PJ. Drug- and estrogen-induced cholestasis through inhibition of the hepatocellular bile salt export pump (Bsep) of rat liver. *Gastroenterology.* 2000;118(2):422-30.
61. Vallejo M, Briz O, Serrano MA, Monte MJ, Marin JJ. Potential role of trans-inhibition of the bile salt export pump by progesterone metabolites in the etiopathogenesis of intrahepatic cholestasis of pregnancy. *J Hepatol.* 2006;44(6):1150-7.
62. Abu-Hayyeh S, Martinez-Becerra P, Sheikh Abdul Kadir SH, Selden C, Romero MR, Rees M, et al. Inhibition of Na⁺-taurocholate Co-transporting polypeptide-mediated bile acid transport by cholestatic sulfated progesterone metabolites. *J Biol Chem.* 2010;285(22):16504-12.
63. Abu-Hayyeh S, Papacleovoulou G, Lovgren-Sandblom A, Tahir M, Oduwole O, Jamaludin NA, et al. Intrahepatic cholestasis of pregnancy levels of sulfated progesterone metabolites inhibit farnesoid X receptor resulting in a cholestatic phenotype. *Hepatology.* 2013;57(2):716-26.
64. Abu-Hayyeh S, Williamson C. Progesterone metabolites as farnesoid X receptor inhibitors. *Dig Dis.* 2015;33(3):300-6.
65. Kullak-Ublick GA, Meier PJ. Mechanisms of cholestasis. *Clin Liver Dis.* 2000;4(2):357-85.
66. Reyes H, Sjoval J. Bile acids and progesterone metabolites in intrahepatic cholestasis of pregnancy. *Ann Med.* 2000;32(2):94-106.
67. Beuers U, Trauner M, Jansen P, Poupon R. New paradigms in the treatment of hepatic cholestasis: from UDCA to FXR, PXR and beyond. *J Hepatol.* 2015;62(1 Suppl):S25-37.
68. Pauli-Magnus C, Stieger B, Meier Y, Kullak-Ublick GA, Meier PJ. Enterohepatic transport of bile salts and genetics of cholestasis. *J Hepatol.* 2005;43(2):342-57.

69. Chojkier M. Troglitazone and liver injury: in search of answers. *Hepatology*. 2005;41(2):237-46.
70. Watkins PB. Insight into hepatotoxicity: The troglitazone experience. *Hepatology*. 2005;41(2):229-30.
71. Watkins PB, Whitcomb RW. Hepatic dysfunction associated with troglitazone. *N Engl J Med*. 1998;338(13):916-7.
72. Yamamoto Y, Yamazaki H, Ikeda T, Watanabe T, Iwabuchi H, Nakajima M, et al. Formation of a novel quinone epoxide metabolite of troglitazone with cytotoxicity to HepG2 cells. *Drug Metab Dispos*. 2002;30(2):155-60.
73. He K, Talaat RE, Pool WF, Reily MD, Reed JE, Bridges AJ, et al. Metabolic activation of troglitazone: identification of a reactive metabolite and mechanisms involved. *Drug Metab Dispos*. 2004;32(6):639-46.
74. Funk C, Ponelle C, Scheuermann G, Pantze M. Cholestatic potential of troglitazone as a possible factor contributing to troglitazone-induced hepatotoxicity: in vivo and in vitro interaction at the canalicular bile salt export pump (Bsep) in the rat. *Mol Pharmacol*. 2001;59(3):627-35.
75. Funk C, Pantze M, Jehle L, Ponelle C, Scheuermann G, Lazendic M, et al. Troglitazone-induced intrahepatic cholestasis by an interference with the hepatobiliary export of bile acids in male and female rats. Correlation with the gender difference in troglitazone sulfate formation and the inhibition of the canalicular bile salt export pump (Bsep) by troglitazone and troglitazone sulfate. *Toxicology*. 2001;167(1):83-98.
76. Yang K, Woodhead JL, Watkins PB, Howell BA, Brouwer KLR. Systems pharmacology modeling predicts delayed presentation and species differences in bile acid-mediated troglitazone hepatotoxicity. *Clin Pharmacol Ther*. 2014;96(5):589-98.
77. Jemnitz K, Veres Z, Vereczkey L. Contribution of high basolateral bile salt efflux to the lack of hepatotoxicity in rat in response to drugs inducing cholestasis in human. *Toxicol Sci*. 2010;115(1):80-8.
78. Fattinger K, Funk C, Pantze M, Weber C, Reichen J, Stieger B, et al. The endothelin antagonist bosentan inhibits the canalicular bile salt export pump: a potential mechanism for hepatic adverse reactions. *Clin Pharmacol Ther*. 2001;69(4):223-31.
79. Leslie EM, Watkins PB, Kim RB, Brouwer KL. Differential inhibition of rat and human Na⁺-dependent taurocholate cotransporting polypeptide (NTCP/SLC10A1) by bosentan: a mechanism for species differences in hepatotoxicity. *J Pharmacol Exp Ther*. 2007;321(3):1170-8.
80. Woodhead JL, Yang K, Siler SQ, Watkins PB, Brouwer KLR, Barton HA, et al. Exploring BSEP inhibition-mediated toxicity with a mechanistic model of drug-induced liver injury. *Front Pharmacol*. 2014;5:240.
81. Ito T, Ozaki Y, Son Y, Nishizawa T, Amuro H, Tanaka A, et al. Combined use of ursodeoxycholic acid and bosentan prevents liver toxicity caused by endothelin receptor antagonist bosentan monotherapy: two case reports. *J Med Case Rep*. 2014;8:250.
82. Hofmann AF. Why bile acid glucuronidation is a minor pathway for conjugation of endogenous bile acids in man. *Hepatology*. 2007;45(4):1083-4; author reply 4-5.
83. Perreault M, Bialek A, Trottier J, Verreault M, Caron P, Milkiewicz P, et al. Role of glucuronidation for hepatic detoxification and urinary elimination of toxic bile acids during biliary obstruction. *PLoS One*. 2013;8(11):e80994.
84. Perreault M, Gauthier-Landry L, Trottier J, Verreault M, Caron P, Finel M, et al. The Human UDP-glucuronosyltransferase UGT2A1 and UGT2A2 enzymes are highly active in bile acid glucuronidation. *Drug Metab Dispos*. 2013;41(9):1616-20.
85. Dawson PA, Karpen SJ. Intestinal transport and metabolism of bile acids. *J Lipid Res*. 2015;56(6):1085-99.
86. Alnouti Y. Bile Acid sulfation: a pathway of bile acid elimination and detoxification. *Toxicol Sci*. 2009;108(2):225-46.

87. Aleo MD, Luo Y, Swiss R, Bonin PD, Potter DM, Will Y. Human drug-induced liver injury severity is highly associated with dual inhibition of liver mitochondrial function and bile salt export pump. *Hepatology*. 2014;60(3):1015-22.
88. Rachek LI, Yuzefovych LV, Ledoux SP, Julie NL, Wilson GL. Troglitazone, but not rosiglitazone, damages mitochondrial DNA and induces mitochondrial dysfunction and cell death in human hepatocytes. *Toxicol Appl Pharmacol*. 2009;240(3):348-54.
89. Allen K, Jaeschke H, Copple BL. Bile acids induce inflammatory genes in hepatocytes: a novel mechanism of inflammation during obstructive cholestasis. *Am J Pathol*. 2011;178(1):175-86.
90. Chen M, Borlak J, Tong W. High lipophilicity and high daily dose of oral medications are associated with significant risk for drug-induced liver injury. *Hepatology*. 2013;58(1):388-96.
91. Schadt S, Simon S, Kustermann S, Boess F, McGinnis C, Brink A, et al. Minimizing DILI risk in drug discovery - A screening tool for drug candidates. *Toxicol In Vitro*. 2015.
92. Aubrecht J, Schomaker SJ, Amacher DE. Emerging hepatotoxicity biomarkers and their potential to improve understanding and management of drug-induced liver injury. *Genome Med*. 2013;5(9):85.
93. Yamazaki M, Miyake M, Sato H, Masutomi N, Tsutsui N, Adam KP, et al. Perturbation of bile acid homeostasis is an early pathogenesis event of drug induced liver injury in rats. *Toxicol Appl Pharmacol*. 2013;268(1):79-89.
94. Coen M, Lenz EM, Nicholson JK, Wilson ID, Pognan F, Lindon JC. An integrated metabonomic investigation of acetaminophen toxicity in the mouse using NMR spectroscopy. *Chem Res Toxicol*. 2003;16(3):295-303.
95. Coen M, Ruepp SU, Lindon JC, Nicholson JK, Pognan F, Lenz EM, et al. Integrated application of transcriptomics and metabonomics yields new insight into the toxicity due to paracetamol in the mouse. *J Pharm Biomed Anal*. 2004;35(1):93-105.
96. Sun J, Ando Y, Ahlbory-Dieker D, Schnackenberg LK, Yang X, Greenhaw J, et al. Systems Biology Investigation to Discover Metabolic Biomarkers of Acetaminophen-Induced Hepatic Injury Using Integrated Transcriptomics and Metabolomics. *J Mol Biomark Diagn*. 2013;S1:002.
97. Luo L, Schomaker S, Houle C, Aubrecht J, Colangelo JL. Evaluation of serum bile acid profiles as biomarkers of liver injury in rodents. *Toxicol Sci*. 2014;137(1):12-25.
98. Luo L, Kenny J, Warner R, Johnson K, Colangelo JL. Profiling individual bile acids in human populations using a UPLC/MS/MS method (abstract). *The Toxicologist: Supplement to Toxicological Sciences*. 2015;144(1):256.
99. Beger RD, Bhattacharyya S, Yang X, Gill PS, Schnackenberg LK, Sun J, et al. Translational biomarkers of acetaminophen-induced acute liver injury. *Arch Toxicol*. 2015.
100. Watanabe T, Miyake M, Shimizu T, Kamezawa M, Masutomi N, Shimura T, et al. Utility of bilirubins and bile acids as endogenous biomarkers for the inhibition of hepatic transporters. *Drug Metab Dispos*. 2015;43(4):459-66.
101. Weiler S, Merz M, Kullak-Ublick GA. Drug-induced liver injury: the dawn of biomarkers? *F1000Prime Rep*. 2015;7:34.